

EMISIÓN DE METANO CAUSADO POR EL GANADO Y EL ESTIÉRCOL EN LA COMARCA LAGUNERA: IMPACTOS AMBIENTALES Y OPORTUNIDADES PARA LA PRODUCCIÓN DE BIOGÁS

METHANE EMISSIONS CAUSED BY CATTLE AND MANURE IN THE LAGUNA REGION: ENVIRONMENTAL IMPACT AND OPORTUNITIES FOR BIOGAS PRODUCTION

M.-Quantin, B. Roy, A. Moreno-Resendez,
J. Vásquez-Arroyo*

Departamento de Agroecología. División de Carreras Agronómicas. UAAAN-UL. Periférico Raúl López Sánchez S/N Torreón, 27000 Coahuila, México. Tel: (871) 729-7645. *Correo-e: Salmonella2004@hotmail.com.

RESUMEN. La Comarca Lagunera representa el 0.03 % del territorio mexicano y cuenta con el 0.2 % de la población lechera nacional, la cual cuenta con aproximadamente 430,000 cabezas de ganado. La industria ganadera produce metano de dos formas. La más importante es la fermentación entérica, derivada de la alimentación de ruminantes, y la segunda es debida al manejo del estiércol. Una vez que el metano es liberado a la atmósfera, se reconoce su importancia como un gas del efecto de invernadero, que es 58 veces más efectivo que el dióxido de carbono. Para estimar la cantidad de metano liberado en esta región, se realizó la estimación de las emisiones de CH₄ originadas por el ganado de la Comarca Lagunera empleando el modelo del Grupo Intergubernamental del Cambio Climático (IPCC). Se determinó la liberación 55 Gg de CH₄ por año, lo cual representa un 2.5 % de las emisiones nacionales de origen ganadero, una gran cantidad considerando el área de la región (0.03 %). El ganado fue responsable del 83 % de la fermentación entérica liberada a la atmósfera, y el restante 17 % se originó de los sistemas de manejo del estiércol. Entre las alternativas para aminorar las emisiones de CH₄, se cuenta con mejorar la calidad de la dieta para una mejor utilización de los insumos proporcionados, por ejemplo, la reducción en la pérdida de energía en las formas de metano y empleo de subproductos del girasol (aceite) o monesina además de raciones basadas en maíz, todo lo cual puede disminuir las emisiones hasta en un 20 %. Sin embargo, posteriores investigaciones requieren precisar este fenómeno y realizar análisis de costo-beneficio y factibilidad a gran escala.

Palabras clave: Metano, biogás, gases de invernadero, cambio climático, establos.

SUMMARY. The Laguna region represents 0.03 % of the Mexican territory and holds 0.2 % of the national milking cow population, with approximately 430,000 heads. Livestock produce methane from two ways. The most important source is enteric fermentation of feed by ruminants, and the second is manure management. Once methane is released to the atmosphere, it is well known to be an important Greenhouse Gas (GHG), 58 times more efficient than carbon dioxide as an Infrared (IR) radiation trap. CH₄ emissions from cattle origin in the Laguna region have been estimated using the IPCC model. Fifty five Gg of CH₄ are produced per year which represents 2.5 % of national CH₄ emissions from livestock origin, a high amount considering the area of the region (0.03 %). For cattle, enteric fermentation is responsible for 83 % of CH₄ released in the atmosphere and 17% comes from manure management systems. Among the existing alternatives for mitigating those CH₄ emissions, improvements in diet quality toward better utilization of ingested energy, i.e. reduction of energy losses in the form of methane, and utilization of supplements such as sunflower oil or monensin in addition to maize based ration may decrease CH₄ emissions up to 20%. However, further investigation is needed to better understand these phenomena and find out cost effective and large scale feasible alternatives.

Keywords. CH₄, biogas, greenhouse gases, climatic change, livestock farming.

INTRODUCTION

The Laguna region, located on the states of Coahuila y Durango in the northern México, is the most important milk production region in the country. This area extends

over 44,887 km², and is located at 101° 40' W Long, 25° 05' N at an altitude of 1,120 meters over the sea level (Anonymous 2006 www.inegi.gob.mx). The Laguna region represents 0.03 % of the Mexican territory and holds 0.2 % of the national milking cow population, with

425,960 heads in 2005 (Anonymous, 2005 www.siap.gob.mx). This cattle is mainly reared in an intensive way in order to supply the needs of an industrialized milk production, such as the LALA or the Bell companies, which are the leaders in the region with a production of about 1.8×10^9 liters/year (Anonymous, 2006 www.siap.gob.mx) mainly intended for exportation out of the Laguna region.

Such a production in a semi arid area is an ecological paradox regarding to the water availability in the region. Indeed, many rivers in the region have been dried due to field irrigation for forage production and the levels of underground water are decreasing dramatically since high yields irrigated crop productions have been started in the Laguna. The region also suffers soil fertility problems on account of maize (*Zea mays*) and alfalfa (*Medicago sativa*) monoculture practices. Sustainability for livestock agroecosystems is thus in at critical point. Cattle rearing and especially large milking cow feedlots are criticized for their contribution to air pollution, i.e. unpleasant smell and methane (CH_4) production. Livestock produce methane from two ways. The most important source is enteric fermentation of feed by ruminants, and the second is manure management (Ruiz-Suárez & Gonzalez-Ávalos 1997), that varies according to storage systems and composting methods at work (Hao *et al.*, 2001; Umetsu *et al.*, 2005). Once methane is released to the atmosphere, it is well known to be an important Greenhouse Gas (GHG), 21 or 58 times more efficient -by molecule or by mass, respectively - than carbon dioxide as an Infrared (IR) radiation trap (Stedd & Hashimoto, 1993). Methane also contributes to the deterioration of the ozone layer (Rhoderick & Dorko 2004). It is the second GHG involved in the global warming (after CO_2 O_3) and its global participation to climate forcing in the past 150 years is 0.57 Wm^{-2} , which is about 35 % of the climate change caused by carbon dioxide (1.6 Wm^{-2}) (Lelieveld *et al.*, 1998). The importance of global warming for environmental stability does not need to be proved nowadays as it has already been the topic of numerous investigations. However, a huge uncertainty remains on its consequences on human and natural ecosystems (Penman *et al.*, 2000).

Methane presence in the atmosphere is also a source of health problems due to its capacity to produce ozone while decomposing in the troposphere. Increase in tropospheric of ozone is associated with premature mortality, respiration y cardiovascular problems, as well as ecosystem perturbations. Mitigation of ozone pollution with better methane emission management brings global health benefits for human societies and the environment (Weast *et al.*, 2006). Methane effects

on global warming and human health raise the problem of the sustainability of methanogenic activities such as cattle rearing. Although many other sectors contribute to methane emissions (industry, transports, etc.), improvements in livestock management - both for enteric fermentation and for manure management - have to be considered seriously. Several ways are available today to influence CH_4 impact on ecosystems, among which are diet improvement toward better utilization of ingested energy (Beachemin & McGinn, 2005) and biogas production. Indeed, methane is an important renewable energy source. Anaerobic digestion (AD) processes for cattle manure composting offer an interesting alternative for farm owners to reduce methane emissions on a local scale while generating bio energy (biogas). Comparing traditional manure management methods AD process presents advantages such as odor elimination, nutrient recovery and mitigation of lixiviation, reduction of pathogen risks and diminution of GHGs emissions (Wilkie, 2005a, 2005b). The Laguna region represents a non negligible potential for biogas production because of its important cattle population, and the availability of both simple y complex biogas production systems gives farmers all the possibilities to get into such a progress.

The objective of this paper is to examine and quantify methane emissions from cattle in the Laguna region, using models given by the Intergovernmental Panel on Climate Change (Houghton *et al.*, 1997; Dong *et al.*, 2006). Further, we will see in more detail how methane impacts on climate and human health, and to explore the existing ways to reduce methane emissions from cattle and finally estimate the biogas potential of the region.

Methane emissions from livestock in the Laguna region

Livestock and especially ruminants (bovine and ovine) produce methane in two ways. The first and most important one is enteric fermentation, where methane originates from carbohydrate reduction by anaerobic micro organisms during the digestion process. CH_4 quantity depends on animal type, age, weight and diet (Carmona *et al.*, 2005). The second way results from anaerobic decomposition of manure, and its intensity increases when an important cattle population is kept in closed areas, i.e. feedlots and "loose-housing" systems (Houghton *et al.*, 1997).

The methane emissions level in a region is an excellent way for identifying mitigation options. Houghton *et al.* (1997) developed a general model to estimate GHGs emissions from anthropogenic sources. The agriculture section presents methods to evaluate methane emissions from cattle, taking in account data such as

cattle population, climate characteristics and manure management practices, and emission factors for both enteric and manure anaerobic fermentation. The IPCC Tier 2 models suffer a lack of precision when applied to the case of Mexico (González-Ávalos & Ruíz-Suárez, 2001). They have determined country-specific emission factors for the national inventory closer to the reality. Those emission factors are used in the present paper for methane emission calculations in the Laguna region. The results show that CH₄ emissions from domestic livestock in the Laguna region are about 55 Gg yr⁻¹ (Table 1), which represent 2.5 % of national emissions, concentrated in 0.03 % of national territory. The biggest part (83 %) is due to enteric fermentation of dairy cattle, i.e. dairy cows (~46 Gg yr⁻¹). The emissions from manure management (17 %), are imputable to poultry farming with ~0.5 Gg yr⁻¹.

The lack of regional data about CH₄ emissions from fuel combustion, industrial processes and waste treatment makes impossible comparison with those sectors, but national data from the national inventory of GHGs 1994-1998 shows that methane from livestock origin is about 25% of the total CH₄ released. Any reduction in this sector would thus have an important impact on these emissions.

Effects of CH₄ on global warming

The emission of CH₄ in the atmosphere presents direct and indirect effects. The direct effect consists in the contribution to climate forcing of the CH₄ molecules. One added CH₄ molecule absorbs infrared radiation about 25 times more efficiently than one added CO₂ molecule (Lelieveld *et al.*, 1993), which can be partly explained by the fact that CO₂ levels are about 200 times higher and therefore its absorption lines are already saturated (Lelieveld *et al.*, 1998). This makes CH₄ being one of the two most important today's greenhouse gas in the atmosphere after CO₂ (with ozone) (Ramaswamy, 2001). Further, decomposition of CH₄ generates formation of other radiatively active gases which increase its direct effect on climate forcing by approximately 30 % (Lelieveld *et al.*, 1998). These gases are tropospheric ozone (O₃), CO₂ and water vapour.

Impacts of CH₄ emissions and of its decomposition products

Importance of greenhouse gases on climate change has not anymore to be demonstrated, as it is commonly accepted as one of most serious impacts of human activities on the environment, both by the scientific community (Houghton *et al.*, 1997) and today, by the wide public. However, besides this primordial aspect, some of the greenhouse gases may procure additional negative effects. It is the case of increasing tropospheric ozone. If global ozone ranks with CH₄ only behind CO₂

as the most important greenhouse gas, its increase in the troposphere implies also other consequences. West *et al.* (2006), identify two main ones. First, premature mortalities can be linked to the high presence of ozone, and especially in the case of cardiovascular and respiratory mortalities. Secondly, increased global ozone affects ecosystems by reducing the net primary productivity of plants. This reduction has for both incidences to diminish plants uptakes of CO₂ - and therefore increasing climate forcing - and to modify agricultural cultures productivity, while changing plants development.

Although CH₄ has greater potential for ozone abatement, it has not been considered for air quality management, contrary to nitrogen oxides (NOx) and nonmethane volatile organic compounds (NMVOCs) (West & Fiore, 2005). Reason for this is that ozone is traditionally considered to be a local and regional problem and that the effects of CH₄ mitigation on surface O₃ are delayed and widespread, offering small local benefits, whereas NOx and NMVOCs affect ozone more rapidly and thus more locally. However, West & Fiore (2005) calculate that a 17 % reduction of current global CH₄ anthropogenic emissions, apart from reducing the radioactive forcing level by ~0.12 W m⁻², would decrease ozone by ~1 ppb. Ozone reduction would prevent about 370 000 all-cause mortalities between 2010 and 2030 and about 30 000 for the year 2030 only. Suppressing about 20 % of the anthropogenic CH₄ emissions is therefore considered as a cost-effective process, as the money saved by avoiding this amount of premature mortalities exceeds the money engaged to implement the CH₄ reduction. Moreover, if this quantity of CH₄ is captured for energy purpose, it would provide ~2 % of current global natural gas production (West *et al.*, 2006). As CH₄ emissions present negative consequences and potential reduction benefits that are globally shared, it seems that their control should be organized at an international level (West & Fiore 2005; West *et al.*, 2006). This does not mean that it prevents to look forward how reduced CH₄ emissions can be implemented at a local scale.

Regarding CO₂ and water vapor emission caused by CH₄ decomposition, it appears that their contribution to the direct radioactive forcing effect of CH₄ increase only about 5 % (Lelieveld *et al.*, 1998). Moreover, the contribution of CO₂ shall not be considered since most of it (~80%) comes from biogenic CH₄, so that it is in fact recycled CO₂ and therefore does not accentuate climate forcing. Concerning water vapor, this one is only relevant when considered in the stratosphere.

CH₄ reaction in the atmosphere

According to Lelieveld *et al.* (1998), CH₄ is mainly removed from its reaction with hydroxyl radicals (OH) in

the atmosphere (-90 %). Hydroxyl radicals, as for them, are formed via the action of solar radiation on ozone and water vapor (R1 y R2), and their removal is almost totally ensured by CO y CH₄.

Reaction between CH₄ and OH leads first to formation of CH₂O and HO₂ (R3-R5). Further, CH₂O degradation generates apparition of carbon monoxide (CO) and then of CO₂ (CO + OH + O₂ → HO₂ + CO₂), while the HO₂ radicals suffer several reactions leading to the formation of O₃ molecules as well as new OH radicals that contribute to renew their presence in the atmosphere (R6-R8). However, this production is dependent on the presence of nitrogen oxides (NO_x). Indeed, when levels of NO_x in the atmosphere are low (less than 25-50 pmol/mol), the hydroperoxy radicals (HO₂) (produced from CH₄ breakdown) cannot react with NO but destroy O₃ and limit the regeneration of OH radicals. Thus, methane oxidation in the atmosphere is a net source of O₃ and OH radicals if NO_x levels are high, whereas net destruction of O₃ and OH prevails if NO_x levels are low. As a consequence of this, the more NO_x in the atmosphere, the more OH and O₃ y the less important is CH₄ lifetime, when knowing that CH₄ lifetime is the quotient of CH₄ burden in the atmosphere and the annual CH₄ breakdown (Lelieveld, 1998).

Reduction pathways

Reducing emissions from enteric fermentation

The most important source of CH₄ emissions from livestock comes from enteric fermentation, it appears primordial to consider alternatives to mitigate the phenomena. According to Carmona *et al.* (2005), methane emissions can vary a lot with the diet submitted to cattle.

Improvements in diet quality can lessen CH₄ emissions by giving to the fermentative process in the rumen better physical chemical conditions and by the way reduce digestive energy loss, which is the cause of methanogenesis. Recent studies in Canada (Beauchemin & McGinn 2005; 2006; McGinn *et al.*, 2004) show that diets supplemented with sunflower oil, ionophores (e.g. monensin), and possibly some yeast products, as well as a corn-based diet during the finishing phase of the production cycle of feedlot cattle can significantly reduce enteric methane production in the rumen, which represent an improvement in feed efficiency. Thus, mitigating CH₄ losses from cattle has both long-term environmental and short-term economic benefits.

Reducing emissions from manure management origin: biogas production

Biogas production from anaerobic digestion of cattle manure appears to be an interesting solution to combine

prevention of global warming, preservation of ozone layer and reduction of health risks due to tropospheric O₃. At the same time it provides bio-energy (methane), as well as a high quality fertilizer for field use. Anaerobic digestion is an engineered methanogenic decomposition of organic matter under oxygen-free conditions (Wilkie, 2005a). The biogas produced in the digester is primarily composed of methane (~60 %) and carbon dioxide (~40 %) (Borole *et al.* 2006), with traces of hydrogen sulphide and ammonia. The quality of the feedstock used in the digester will determine how much gas is produced (British-Biogen, 2000).

The CH₄ generated can be used as a fuel for an engine connected to a generator to produce electricity that can be used on site or marketed to the network. The latter possibility is by far the most interesting one for big scale anaerobic digestion units as it leads to an additional net income for the farm owner. Biogas can also be used to produce heat or merely be burned for kitchen use in the case of small-scale farm production (Laichena & Wafula, 1997). The electricity generated can cover totally or partially the needs of farm machinery such as milking or maintaining machine, depending on the farm size.

In contrast with classical storage and aerobic composting techniques, whose methane production escape in the atmosphere and suffers a loss of nutrients (nitrogen) in the form of gas or by lixiviation (Hao *et al.*, 2001; Umetsu *et al.*, 2005), anaerobic systems save those products inside the closed system of the biodigester and provide them as mineralized and more soluble nutrients for field use (Wilkie, 2005).

Moreover, anaerobic digestion can reduce the potential for global warming in two ways. First, the use of biogas as an energy source replaces the consumption of fossil fuels such as coal, oil and natural gas, and thus CO₂ emissions related to that combustion is avoided. Indeed, CO₂ emitted from biogas burning comes from carbon of cattle feed, which originates from photosynthesis processes in feed or forage production, and is part of a closed carbon cycle and, therefore, does not contribute to increasing atmospheric CO₂ levels. The second way is that anaerobic digestion reduces CH₄ release in the atmosphere and by the way mitigates potential global warming, as CH₄ is 21 times more efficient as CO₂ for climate forcing (Wilkie, 2005a).

Furthermore, CH₄ reduction also influences O₃ production in the troposphere to prevent premature mortality and other diseases at a local scale (Lelieveld *et al.* 1998).

Biogas potential in the Laguna region for electricity production

The cattle population, for the Laguna region represents a great potential for energy production. Electricity

produced from biogas has an average yield of 30 % as the biggest part of the energy is converted by the generator into heat. Nevertheless, this heat can be recovered to warm water or directly to the digester to improve its performance (British-Biogen, 2000). Taking in account cattle population, average slurry and biogas production, we can estimate the electricity potential of the Laguna region. If all the slurry produced were digested by AD process and converted to electricity, it would generate around 61 GW h⁻¹. It is evident that this level of electricity production will never be reached because all farms are not going to set up biogas plants in the next few years, but many projects have already begun in the Laguna region with the support of the United Nations and Ag Cert Company. A project of installing 28 biodigestors in the region in the next two months, totally financed by the United Nations, is ongoing in collaboration with big farm owners (more than 700 cows). On the other hand, the Universidad Autónoma Agraria "Antonio Narro" (UAAAN) is working on biogas application to small scale farms and rural communities in order to bring this technology to the countryside and give to rural community the opportunity to produce their own energy and by self sufficient.

DISCUSSION

CH₄ emissions from cattle origin in the Laguna region have been estimated using the IPCC model (Houghton *et al.*, 1997). Fifty five Gg of CH₄ are produced per year which represents 2.5 % of national CH₄ emissions from livestock origin, a high amount considering the area of the region (0.03 %). Indeed, cattle rearing in the region are large and suffer serious sustainability problems. For cattle, enteric fermentation is responsible for 83 % of CH₄ released in to the atmosphere and 17 % comes from manure management systems. The greenhouse potential of methane as well as the negative effects of its decomposition by-products on human health and air pollution turns CH₄ into a threat for global climate balance and natural and human ecosystems stability. As CH₄ emissions from livestock is about 25 % of total CH₄ emissions, any reduction on this sector would be environmentally effective and a path toward more sustainability and consciousness of our surroundings. Among the existing alternatives for mitigating those CH₄ emissions, improvements in diet quality toward better utilization of ingested energy, i.e. reduction of energy losses in the form of methane, and utilization of supplements such as sunflower oil or monensin in addition to maize based ration may decrease CH₄ emissions up to 20 %. However, further investigation is needed to better understand these phenomena and find out cost effective and large scale feasible alternatives. Considering the manure management issue, utilization of AD technology to control manure fermentation may

avoid the majority of CH₄ emissions due to manure decomposition and of the same time generate a renewable and environmentally friendly energy: biogas. Utilization of biogas on farms can reduce or eliminate electricity consumption due to farming activity. Digesting animal slurry makes the farming system more sustainable and self sufficient, and helps farmers and rural communities being aware of their impact on the environment and of the power they have to convert this problem into damage-free activities.

In the case of the Laguna region, if a process carries several advantages but, one should not forget that the main problem still remains: the high concentration of animals for intensive production. Utilization of AD technologies and especially the biogas production make it possible to diminish some negative effects of high concentrations of domestic animals, but do not represent a solution to the whole problem. Dramatic decrease in water availability, high level of soil erosion and widening social gaps are some of the actual indicators that show that the presence of intensive production in the region is not compatible with the wish to reach sustainable and long-term development. AD technologies represent interesting tools for future development only if considered as a part of the greater and radical changes that have to be implemented in the agricultural sector and in the society in general.

Acknowledges.

We acknowledges to the project support by Universidad Autónoma Agraria "Antonio Narro" with register number 02-03-1510-2798.

REFERENCES

- Anonymous. 2006. Instituto Nacional de Estadística, Geografía e Informática. Disponible en: www.inegi.gob.mx.
- Anonymous. 2005. Servicio de Información Agroalimentaria and Pesquera. Disponible en www.siap.gob.mx.
- Beauchemin KA, y McGinn SM. 2005. Methane emissions from feedlot cattle fed barley or corn diets. *J Anim Sci* 83:653-61.
- Beauchemin KA, y McGinn SM. 2006. Methane emissions from beef cattle: effects of fumaric acid, essential oil, and canola oil. *J Anim Sci* 84:1489-96.
- Borole AP, Klasson KT, Ridenour W, Holland J, Karim K, y Al-Dahhan MH. 2006. Methane production in a 100-L upflow bioreactor by anaerobic digestion of farm waste. *Appl Biochem Biotechnol* 129-132:887-96.
- British-Biogen. 2000. Anaerobic digestion farm and food processing residues: good practice guidelines. Available in: <http://www.britishbiogen.co.uk/gpg/adgpg/adgpg.pdf>.
- Carmona JC, Bolivar DM, y Giraldo LA. 2005. El gas metano en la producción ganadera and alternativas para

- medir sus emisiones and aminorar su impacto a nivel ambiental and productivo. *Rev Col Cienc Pec* 18:49-63.
- Dong H, Mangino J, McAllister TA, Hatfield JL, Johnson DE, Lassey KR, Aparecida de Lima M, Romanovsky A, Bartram D, Gibb D, y Martin Jr. JH. 2006. Emission from livestock and manure management. In S Eggleston, L Buendia, K Miwa, T Ngara and K Tanabe (eds.): 2006. IPCC guideline for national greenhouse gas inventories. Agriculture, forestry and other land use. Japan: IGES, pp. 10.8-10.87.
- González-Avalos E, y Ruiz-Suárez JC. 2007. Methane conversion factors from cattles manure in México. *Atmosfera* 20:83-92.
- Gonzalez-Avalos E, y Ruiz-Suarez LG. 2001. Methane emission factors from cattle manure in Mexico. *Bioresour Technol* 80:63-71.
- Hao X, Chang C, Larney FJ, y Travis GR. 2001. Greenhouse gas emissions during cattle feedlot manure composting. *J Environ Qual* 30:376-86.
- Houghton JT, Meira-Fio LG, Kim B, Treaton K, Mamaty I, Bonduki A D, Griggs DJ, y Callender BA. 1997. IPCC Guidelines for National Greenhouse Gas Inventories. Bracknell, UK: IPCC WGI.
- Laichena JK, y Wafula JC. 1997. Biogas technology for rural households in kenya. *OPEC Rev* 21:223-224.
- Lelieveld J, Crutzen PJ, y Brühl C. 1993. Climate effects of atmospheric methane. *Chemosphere* 26:739-768.
- Lelieveld J, Crutzen PJ, y Dentener FJ. 1998. Changing concentration, lifetime and climate forcing of atmospheric methane. *Tellus* 50B:128-150.
- McGinn SM, Beauchemin KA, Coates T y Colombatto D. 2004. Methane emissions from beef cattle: Effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *J Anim Sci* 82:3346-56.
- Penman J, Kruger D, Galbally I, Hiraishi T, Nyenzi B, Emmanuel S, Buendia L, Hoppaus R, Martinsen T, Meijer J, Miwa K y Tanabe K. 2000. Good practice guidance and uncertainty management in national greenhouse gas inventories. Tokio, Japan: Institute for Global Strategy.
- Ramaswany V, Boucher O, Haigh J, Hauglustaine D, Haywood J, Myhre G, Nakajima T, Shi GY, y Solomon S. 2001. The climatic basis. In JT Houghton, AND Ding, DJ Griggs, M Noguer, PJ Van der Linden, X Dai, K Markell y CA Johnson (eds.): Climatic change. Cambridge, U.K.: Cambridge University Press, pp. 349-416.
- Rhoderick GC, y Dorko WD. 2004. Standar development of global warming gas species: methane, nitrous oxide, trichlorofluoromethene, and dichlorodifluoromethene. *Environ Sci Technol* 38:2685-92.
- Ruiz-Suarez LG, y Gonzalez-Avalos E. 1997. Modeling methane emissions from cattle in Mexico. *Sci Total Environ* 206:177-86.
- Steed Jr J, and Hashimoto AG. 1993. Methane emission emission factors for typical US livestock manure management systems. Oregon, USA: Oregon State University, pp. 4.
- Umetsu K, Kimura AND, Takhashi J, Kishimoto T, Kojima T, y Young B. 2005. Methane emission from stored dairy manure slurry and slurry after digestion by methane digester. *J Anim Sci* 76:73-79.
- West JJ, y Fiore AM. 2005. Management of tropospheric ozone by reducing methane emissions. *Environ Sci Technol* 39:4685-91.
- West JJ, Fiore AM, Horowitz L y Mauzerall DL. 2006. Global health benefits of mitigating ozone pollution with methane emission controls. *Proc Natl Acad Sci USA* 103:3988-3993.
- Wilkie AC. 2005. Anaerobic digestion: Biology and benefits: Natural Resource, Agriculture, and Engineering Service. Ithaca, New York: Cornell University, pp. 63-72.
- Wilkie AC. 2005a. Animal manure management: Innovative technology for odor control, energy production and nutrient recovery. Gainesville, FL, USA: University of Florida.